

## Answer on Question #49930 – Chemistry – Organic Chemistry

### Question:

How we decide which reducing or oxidizing agent is strong and weak and which we should use for the reaction.

### Answer:

We can use the value of measured standard potentials for a wide variety of chemical substances, some of which are listed in Table 1 "Standard Potentials for Selected Reduction Half-Reactions at 25°C". These data allow us to compare the oxidative and reductive strengths of a variety of substances. The half-reaction for the standard hydrogen electrode (SHE) lies more than halfway down the list in Table 1 "Standard Potentials for Selected Reduction Half-Reactions at 25°C". All reactants that lie *above* the SHE in the table are stronger oxidants than H<sup>+</sup>, and all those that lie below the SHE are weaker. The strongest oxidant in the table is F<sub>2</sub>, with a standard electrode potential of 2.87 V. This high value is consistent with the high electronegativity of fluorine and tells us that fluorine has a stronger tendency to accept electrons (it is a stronger oxidant) than any other element.

Table 1. Standard Potentials for Selected Reduction Half-Reactions at 25°C

Half-Reaction	E° (V)
$F_2(g) + 2e^- \rightarrow 2F^-(aq)$	2.87
$H_2O_2(aq) + 2H^+(aq) + 2e^- \rightarrow 2H_2O(l)$	1.78
$Ce^{4+}(aq) + e^- \rightarrow Ce^{3+}(aq)$	1.72
$PbO_2(s) + HSO_4^-(aq) + 3H^+(aq) + 2e^- \rightarrow PbSO_4(s) + 2H_2O(l)$	1.69
$Cl_2(g) + 2e^- \rightarrow 2Cl^-(aq)$	1.36
$Cr_2O_7^{2-}(aq) + 14H^+(aq) + 6e^- \rightarrow 2Cr^{3+}(aq) + 7H_2O(l)$	1.23
$O_2(g) + 4H^+(aq) + 4e^- \rightarrow 2H_2O(l)$	1.23
$MnO_2(s) + 4H^+(aq) + 2e^- \rightarrow Mn^{2+}(aq) + 2H_2O(l)$	1.22
$Br_2(aq) + 2e^- \rightarrow 2Br^-(aq)$	1.09
$NO_3^-(aq) + 3H^+(aq) + 2e^- \rightarrow HNO_2(aq) + H_2O(l)$	0.93
$Ag^+(aq) + e^- \rightarrow Ag(s)$	0.80
$Fe^{3+}(aq) + e^- \rightarrow Fe^{2+}(aq)$	0.77
$H_2SeO_3(aq) + 4H^+ + 4e^- \rightarrow Se(s) + 3H_2O(l)$	0.74

Half-Reaction	$E^\circ$ (V)
$O_2(g) + 2H^+(aq) + 2e^- \rightarrow H_2O_2(aq)$	0.70
$MnO_4^-(aq) + 2H_2O(l) + 3e^- \rightarrow MnO_2(s) + 4OH^-(aq)$	0.60
$MnO_4^{2-}(aq) + 2H_2O(l) + 2e^- \rightarrow MnO_2(s) + 4OH^-(aq)$	0.60
$I_2(s) + 2e^- \rightarrow 2I^-(aq)$	0.54
$H_2SO_3(aq) + 4H^+(aq) + 4e^- \rightarrow S(s) + 3H_2O(l)$	0.45
$O_2(g) + 2H_2O(l) + 4e^- \rightarrow 4OH^-(aq)$	0.40
$Cu^{2+}(aq) + 2e^- \rightarrow Cu(s)$	0.34
$AgCl(s) + e^- \rightarrow Ag(s) + Cl^-(aq)$	0.22
$Cu^{2+}(aq) + e^- \rightarrow Cu^+(aq)$	0.15
$Sn^{4+}(aq) + 2e^- \rightarrow Sn^{2+}(aq)$	0.15
$2H^+(aq) + 2e^- \rightarrow H_2(g)$	0.00
$Sn^{2+}(aq) + 2e^- \rightarrow Sn(s)$	-0.14
$2SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightarrow S_2O_6^{2-}(aq) + 2H_2O(l)$	-0.22
$Ni^{2+}(aq) + 2e^- \rightarrow Ni(s)$	-0.26
$PbSO_4(s) + 2e^- \rightarrow Pb(s) + SO_4^{2-}(aq)$	-0.36
$Cd^{2+}(aq) + 2e^- \rightarrow Cd(s)$	-0.40
$Cr^{3+}(aq) + e^- \rightarrow Cr^{2+}(aq)$	-0.41
$Fe^{2+}(aq) + 2e^- \rightarrow Fe(s)$	-0.45
$Ag_2S(s) + 2e^- \rightarrow 2Ag(s) + S^{2-}(aq)$	-0.69
$Zn^{2+}(aq) + 2e^- \rightarrow Zn(s)$	-0.76
$Al^{3+}(aq) + 3e^- \rightarrow Al(s)$	-1.662
$Be^{2+}(aq) + 2e^- \rightarrow Be(s)$	-1.85
$Li^+(aq) + e^- \rightarrow Li(s)$	-3.04