

## Answer on Question#53117 – Physics / Mechanics – Kinematics – Dynamics

4.905 kJ

### Question

A cubicle vessel of height 1 m is full of water. What will be the minimum work done in taking water out from vessel?

### Solution

Consider an infinitely small volume  $dV$  of water (elementary volume).

Minimum work to take it out from vessel – transfer to the highest point of the vessel ( $dW = g(h - h_i)dm$ ) and shift horizontally out of vessel (assume that there is no friction with medium (e.g air), so no contribution to work there).

$g$  – gravitational acceleration,  $h$  – height of vessel,  $h_i$  – initial height of elementary volume,  $dm$  – mass.

Thus, our aim to calculate work over all heights  $h_i$ . Assume that the lowest point has  $h_i = 0$  m, accordingly to the data, the highest point has  $h_i = 1$  m then (Note: no effect on result). At the same time (in general) we can rewrite  $dm = \frac{dm}{dh}dh$ , and due to assumption that density of water independent from height, we can replace  $\frac{dm}{dh}$  with  $\frac{m}{h}$ . Finally, we can jot down proper integral:

$$W = \int_{h_{imin}}^{h_{imax}} g(h - h_i) \frac{m}{h} dh_i$$

Let us put in actual boundaries and solve it:

$$\begin{aligned} W &= \int_0^1 g(h - h_i) \frac{m}{h} dh_i = \frac{mg}{h} \int_0^1 (h - h_i) dh_i = mg \int_0^1 dh_i - \frac{mg}{h} \int_0^1 h_i dh_i = \\ &= mgh_i \Big|_0^1 - \frac{mg}{h} \frac{h_i^2}{2} \Big|_0^1 = mg - \frac{mg}{2h} = mg \left( 1 - \frac{1}{2h} \right) \end{aligned}$$

There are few numbers from tables we required to know to get numeric answer: (mass of 1  $m^3$  of water)  $m \approx 1000$  kg, (gravitational acceleration)  $g \approx 9.81$   $ms^{-2}$ . Substitute them:

$$W = mg \left( 1 - \frac{1}{2h} \right) = 1000 * 9.81 \left( 1 - \frac{1}{2 * 1} \right) = 4905 J = 4.905 kJ$$